



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by the PROPOSER.

Using a figure similar to the one used in problem 116, let $AOD=\theta_1$, $BOD=\theta_2$, $COD=\theta_3$, etc. Let p be the required chance, p_1 , the chance that it does not contain the center of the circle.

$$p_1 = \frac{2 \int_0^{\frac{1}{2}\pi} \int_0^{\theta_1} \int_0^{\theta_2} \dots \int_0^{\theta_{n-2}} d\theta_1 d\theta_2 d\theta_3 \dots d\theta_{n-1}}{\int_0^{\pi} \int_0^{\theta_1} \int_0^{\theta_2} \dots \int_0^{\theta_{n-2}} d\theta_1 d\theta_2 d\theta_3 \dots d\theta_{n-1}} = \frac{(n-1)! \cdot 2(\frac{1}{2}\pi)^{n-1}}{\pi^{n-1} \cdot (n-1)!} = \frac{1}{2^{n-2}}.$$

$$p=1-p_1=1-(1/2^{n-2}).$$

Also solved with the same result by J. SCHEFFER.

MISCELLANEOUS.

133. Proposed by HARRY S. VANDIVER, Bala. Pa.

If a group G of order mn has a subgroup H of order n , and if n has no prime factor which is less than m , show that H must be a self-conjugate subgroup. (Frobenius.)

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let the substitution of the given subgroup (H) be $1, t_2, t_3, \dots, t_p$.

If H is not self-conjugate, multiply it into any substitution (S) which is not commutative to it. If the transform (H_1) of (H) with respect to any one of these products contains a substitution ($s^{-1}ts$) of order m^β ($\beta > 0$) which is not found in H and if t does not transform H_1 into itself we can form the following rectangle with m conjugate rows of p elements:

$$\begin{array}{ccccccc} s, & st_2, & st_3, & \dots, & st_p & & \\ ts, & tst_2, & tst_3, & \dots, & tst_p & & \\ st^2, & t_2st^2, & t_3st^2, & \dots, & t_pst^2 & & \\ \dots & \dots & \dots & \dots & \dots & & \\ st^m-1, & t_2st^m-1, & t_3st^m-1, & \dots, & t_pst^m-1. & & \end{array}$$

All the substitutions of a given row transform H into the same group, while any two substitutions from different rows transform H into two different groups. As no substitutions in the given rectangle can be equal to each other, the entire group would have to contain at least $p(m+1)$ substitutions. This is contrary to the hypothesis. Therefore H is a self-conjugate sub-group.

134. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

Give a complete solution of the Jacobian equation $\kappa^2 \sin^4 u + 2\kappa^2 \sin^2 u + 1 = 0$.